


## Cellular Potts models

### Differential Adhesion: cell sorting in two dimensions



Yellow cells engulf the red cells as a result of differential adhesion



#### Introduction

This model shows the original cellular Potts model (a.k.a. Glazier-Graner model) of cell sorting based on the Steinberg's differential adhesion hypothesis.

#### Model description

Two `CellTypes` are defined, each of which has a `VolumeConstraint` specifying the cell's target area/volume. In the CPM element, the `MetropolisKinetics` can be configured and the Interaction energies between cell types are specified.

Although cells can be initialized as single points using e.g. the `InitCircle` plugin, in this example, the `Nodes` of each `Cell` in the `CellPopulations` are given explicitly. In fact, these `Populations` are restored results of a previous simulation.

The simulation shows two populations of spatially resolved cells that initially organized in a mosaic fashion. Through differential adhesion, the motile cells sort out and re-organize into an distribution in which one cell type engulfs the other.

Snapshots of the simulation are saved to files named `[Title][Time].xml.gz`. These files containing intermediate and result states can be opened and used as initial conditions for new simulations. Remember to change `StartTime` and `StopTime` accordingly.

## Model

h CellSorting\_2D.xml |h

```
extern>http://imc.zih.tu-dresden.de/morpheus/examples/CPM/CellSorting_2D.xml
```


In Morpheus GUI: File → Examples → CPM → CellSorting\_2D.xml


## Reference

Graner F, Glazier J. [Simulation of Biological Cell Sorting Using a Two-Dimensional Extended Potts Model](#). *Phys. Rev. Lett.* 69 (13): 2013–2016, 1992.

---

## Proliferation in two dimensions



 Growing cell population



## Introduction

This model show a simulation of a growing cell population, using the cellular Potts model.

## Model description

This model specifies CellType which has a VolumeConstraint and a Proliferation plugin.

In the Proliferation plugin, the Conditions for a cell to divide are given. Here, each cell that has more than 90% of the target volume has a small probability to divide. Once a division has taken place, the Equation defined in the Triggers elements are triggered.

## Model


h Proliferation\_2D.xml |h

```
extern>http://imc.zih.tu-dresden.de/morpheus/examples/CPM/Proliferation_2D.xml
```

In Morpheus GUI: File → Examples → CPM → Proliferation\_2D.xml

---

## Persistence



Persistence of individual cells causes self-organized collective motion



## Introduction

This example shows self-organized collective motion of cells as a result of persistence ('cellular inertia'). A similar model has recently been used in ([Czirok et al., 2013](#)).

## Description

The model uses the Persistence plugin that causes cells to prefer to move in their current direction. The direction is stored in a PropertyVector that is used to plot the color and arrows in Gnuplotter.

The model is simulated in a circular domain with constant boundary conditinos, which can be set up in Lattice / Domain / Circle. The value for the constant boundary is specified in CPM / BoundaryValue.

### Try it!

- Change the boundary conditions from circular=constant to x/y=periodic and observe the resulting collective motion.
- Change the decay - time of Persistence (specifying the 'memory').

### Model

h Persistence\_2D.xml |h


```
extern>http://imc.zih.tu-dresden.de/morpheus/examples/CPM/Persistence_2D.xml
```

### Reference

Czirók A, Varga K, Mehes E, Szabo A, [Collective cell streams in epithelial monolayers depend on cell adhesion](#). *New J. Phys.* 15 075006, 2013.

---

### Proliferation in three dimensions



 Cell population grown from single initial cell



## Introruction

This model show a CPM simulation of a growing cell population in 3D.

## Model description

This model specifies `CellType` which has a `VolumeConstraint` and a `Proliferation` plugin. In the `Proliferation` plugin, the `Conditions` for a cell to divide are given. Here, each cell that has more than 90% of the target volume has a small probability to divide. Once a division has taken place, the `Equation` defined in the `Triggers` elements are triggered.

In this model, two medium cell types have been defined. One of these (called `matrix`) is used to represent a matrix with higher adhesivity. This is done by (1) defining the '`matrix`' cell type as a `BoundaryCondition` of the `-z` boundary in the CPM and (2) providing lower contact energy for cell-matrix interaction than for cell-medium interactions.

The simulation is visualized using the `TiffPlotter` that saves TIFF image stacks that can be loaded by image analysis software such as [Fiji](#) and displayed using Fiji's 3D Viewer plugin.

## Model

h Proliferation\_3D.xml |h

```
extern>http://imc.zih.tu-dresden.de/morpheus/examples/CPM/Proliferation_3D.xml
```

In Morpheus GUI: File → Examples → CPM → Proliferation\_3D.xml

Last update: 00:51 30.07.2013 examples:cellular\_potts [https://imc.zih.tu-dresden.de/wiki/morpheus/doku.php?id=examples:cellular\\_potts&rev=1375138306](https://imc.zih.tu-dresden.de/wiki/morpheus/doku.php?id=examples:cellular_potts&rev=1375138306)

---

From: <https://imc.zih.tu-dresden.de/wiki/morpheus/> - **Morpheus**

Permanent link: [https://imc.zih.tu-dresden.de/wiki/morpheus/doku.php?id=examples:cellular\\_potts&rev=1375138306](https://imc.zih.tu-dresden.de/wiki/morpheus/doku.php?id=examples:cellular_potts&rev=1375138306)

Last update: **00:51 30.07.2013**

