



# Multiscale models

## MembraneProperties: Cell polarization and chemotaxis

*Note: MembraneProperties are not available in public version of Morpheus.*



Cell dynamically re-polarizes in response to switching external gradient



### Introduction

This model of cell polarity shows the coupling of three model formalisms:

- A cellular Potts model
- A PDE model, solved on the membrane of the cell
- And an external gradient.

The cell membrane polarizes in response to the external gradient. Chemotactic cell motility depends on the polarity of the cell and the external gradient.

### Description

This example implements two models of cell polarity: Meinhardt's substrate-depletion model and Edelstein-Keshet's wave-pinning model. The user can switch polarity model by Disabling/Enabling the relevant System.

The model defines a one-dimensional reaction-diffusion system (MembraneProperty) representing membrane-bound molecules, and is mapped to a cellular Potts model defining a discrete shaped cell. An external gradient, specified in a PDE, provides a signal for the polarization of the cell. In turn, the polarity of the cell influences its chemotactic behavior.

After a switch in direction of the gradient, the cell re-polarizes in the new direction and starts to move up the gradient, iff the wave pinning model has been selected.

## Model

h CellPolarity.xml |h

```
extern>  
http://imc.zih.tu-dresden.de/morpheus/examples/Multiscale/CellPolarity.xml
```

*Note: This model is not available in Morpheus GUI.*

From:  
<https://imc.zih.tu-dresden.de/wiki/morpheus/> - **Morpheus**

Permanent link:  
<https://imc.zih.tu-dresden.de/wiki/morpheus/doku.php?id=examples:multiscale&rev=1629299761>

Last update: **17:16 18.08.2021**

